



QUEEN'S UNIVERSITY DEPARTMENT OF ANESTHESIOLOGY

Subject: Perioperative Management of Monoamine Oxidase Inhibitors (MAOI)

NUMBER	
PAGE	1 of 2
ORIGINAL ISSUE	12.2000
REVIEWED	10.2003
REVISION	10.2003

Trade Name

Parnate
Nardil
Manerix

Generic Name

Tranlycypromine
Phenelzine
Moclobemide

Introduction:

The classic MAOI, phenelzine and tranlycypromine, irreversibly inhibit MAO for 2 to 3 weeks until new enzyme is synthesized.¹ Moclobemide, a reversible inhibitor of MAO-A (RIMA) causes enzyme inhibition for less than 24 hours.^{2,3} Due to potential drug interactions, discontinuation of classic MAOI 2 to 3 weeks before surgery has been recommended¹ and suitability of these patients for ambulatory anesthesia has been controversial.

Reactions to meperidine (including agitation, hyper/hypotension, convulsions, hyperthermia, and coma.³) have been described for all MAOI^{1,4,5} and RIMA^{2,3} medications. Cocaine also may cause excitatory reactions.⁶ Patients may have an exaggerated response to indirect-acting sympathomimetics as long as 3 weeks after classic MAOI are discontinued.¹ Regarding RIMA's, a patient taking moclobemide had a normal response to both phenylephrine and ephedrine after omitting the morning dose.² Vasopressors at concentrations contained in local anesthetics are not likely to be significantly potentiated in otherwise healthy patients on an MAOI.¹¹

Multiple case reports describe using a variety of general and regional anesthetic techniques for patients on MAOIs without complications⁷⁻¹⁵. Withdrawal of MAOI 2 weeks prior to elective surgery is not without risks and acute exacerbation of depression with suicidal ideation has been reported.¹⁷

Although adverse drug interactions are possible and have been reported in a minority of patients on MAOI, a wide range of anesthetic agents

has been described that permit the safe administration of anesthesia. Therefore, patients taking either classic or selective MAOI are suitable for ambulatory anesthesia without discontinuing their MAOI preoperatively if careful attention is paid to their anesthetic management.

Policy:

1. Patients taking MAOIs may continue taking their medication throughout the perioperative period.
2. Meperidine, cocaine and indirect-acting catecholamines should be avoided in these patients.

References:

1. Wells DG, Bjorksten AR. Monoamine oxidase inhibitors revisited. *Can J Anaesth* 1989; 36: 64-74.
2. Martyr JW, Orlikowski CEP. Epidural anaesthesia, ephedrine and phenylephrine in a patient taking moclobemide, a new monoamine oxidase inhibitor. *Anaesthesia* 1996; 51: 1150-2.
3. McFarlane HJ. Anaesthesia and the new generation monoamine oxidase inhibitors. *Anaesthesia* 1994; 49: 597-9.
4. Stack CG, Rogers P, Linter SPK. Monoamine oxidase inhibitors and anaesthesia. A review. *Br J Anaesth* 1988; 60: 222-7.
5. Evans-Prosser CDG. The use of pethidine and morphine in the presence of monoamine oxidase inhibitors. *Br J Anaesth* 1968; 40: 279-82.
6. Tordoff SG, Stubbing JF, Linter SPK. Delayed excitatory reaction following interaction of cocaine and monoamine oxidase inhibitor (phenelzine). *Br J Anaesth* 1991; 66: 516-8.
7. El-Ganzouri AR, Ivankovich AD, Braverman B, McCarthy R. Monoamine oxidase inhibitors: should they be discontinued preoperatively? *Anesth Analg* 1985; 64: 592-
8. O'Hara JF, Maurer WG, Smith MP. Sufentanil-isoflurane-nitrous oxide anesthesia for a patient treated with monoamine oxidase inhibitor and tricyclic antidepressant. *J Clin Anesth* 1995; 7: 148 -50.
9. Ebrahim ZY, O'Hara J, Borden L, Tetzlaff J. Monoamine oxidase inhibitors and elective surgery. *Cleve Clin J Med* 1993; 60: 129-30.
10. Errando CL, Mateo E Lopez-Alarcon D, Moliner S. Severe interactions with classic and selective monoamine oxidase inhibitors. *Can J Anaesth* 1998; 45: 706-9.
11. Boakes AJ, Laurence DR, Teoh PC et al. Interactions between sympathomimetic amines and antidepressant agents in man. *BMJ* 1973; 1: 311-5.
12. Fischer SP, Mantin R, Brock-Utne JG. Ketorolac and propofol anesthesia in a patient taking chronic monoamine oxidase inhibitors. *J Clin Anesth* 1996; 8: 245-7.

13. Powell H. Use of alfentanil in a patient receiving monoamine oxidase inhibitor therapy. *Br J Anaesth* 1990; 64: 528-9.
14. Doyle DJ. Ketamine induction and monoamine oxidase inhibitors. *J Clin Anesth* 1990; 2: 324-5.
15. Ure DS, Gillies MA, James KS. Safe use of remifentanil in a patient treated with the monoamine oxidase inhibitor phenelzine. *Br J Anaesth* 2000; 84: 414-6.
16. Sides CA. Hypertension during anaesthesia with monoamine oxidase inhibitors. *Anaesthesia* 1987; 42: 633-5.
17. Abdi S, Fishman SM. Acute exacerbation of depression after discontinuation of monoamine oxidase inhibitor prior to cardiac surgery. *Anesth Analg* 1996; 83: 656-7.